

OTH\$ Pre-AP Chemistry Equations and Constants

The following symbols have the definitions specified unless otherwise noted.

g	= gram(s)
nm	= nanometer(s)
atm	= atmosphere(s)
L, mL	= liter(s), milliliter(s)

J, kJ	= joule(s), kilojoule(s)
mol	= mole(s)
mm Hg	= millimeters of mercury
Pa, kPa	= pascal(s), kilopascal(s)

ATOMS, ENERGY, & ELECTRONS

$$E_{\text{photon}} = h\nu$$

$$c = \lambda\nu$$

$$E_{\text{photon}} = \frac{hc}{\lambda}$$

E = energy

ν = frequency

λ = wavelength

Planck's constant, h = $6.626 \times 10^{-34} \text{ J} \cdot \text{s}$

Speed of light, c = $3.00 \times 10^8 \text{ m} \cdot \text{s}^{-1}$

Avogadro's number = $6.02 \times 10^{23} \text{ mol}^{-1}$

ACIDS and BASES

$$K_w = 1.0 \times 10^{-14} \text{ @ } 25^\circ\text{C}$$

$$K_w = [H^+][OH^-]$$

$$pH = -\log[H^+], \quad pOH = -\log[OH^-]$$

$$[H^+] = 10^{-pH}, \quad [OH^-] = 10^{-pOH}$$

$$pH + pOH = 14.00$$

OTHER FORMULAE

$$\text{Percent error} = \left| \frac{\text{accepted value} - \text{experimental value}}{\text{accepted value}} \right| \times 100$$

$$\text{Percent Yield} = \left(\frac{\text{actual yield}}{\text{theoretical yield}} \right) \times 100$$

GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_{\text{total}} = P_1 + P_2 + P_3 \dots$$

$$\frac{P_1V_1}{T_1n_1} = \frac{P_2V_2}{T_2n_2}$$

$$K = ^\circ\text{C} + 273.15$$

$$n = \frac{m}{\text{molar mass}}$$

$$\text{Kinetic Energy} = \frac{1}{2}mv^2$$

$$\text{Molarity, } M = \frac{\text{moles solute}}{\text{liters solution}}$$

$$\text{Dilution formula, } M_1V_1 = M_2V_2$$

$$D = \frac{m}{V}$$

Temperature Units:

K = Kelvin temperature $^\circ\text{C}$ = Celsius temperature

P = pressure

V = volume

T = temperature

n = number of moles of gas

R = Ideal Gas Constant = $8.314 \text{ L} \cdot \text{kPa} \cdot \text{mol}^{-1}\text{K}^{-1}$

= $0.08206 \text{ L} \cdot \text{atm} \cdot \text{mol}^{-1}\text{K}^{-1}$

= $62.36 \text{ L} \cdot \text{torr} \cdot \text{mol}^{-1}\text{K}^{-1}$

= $62.36 \text{ L} \cdot \text{mmHg} \cdot \text{mol}^{-1}\text{K}^{-1}$

= $8.314 \text{ J} \cdot \text{mol}^{-1}\text{K}^{-1}$

$1.000 \text{ atm} = 760.0 \text{ mm Hg} = 760.0 \text{ torr} = 101.3 \text{ kPa}$

STP = 273.15 K and 1.000 atm

Volume of an Ideal Gas @ STP = $22.4 \text{ L} \cdot \text{mol}^{-1}$

v = velocity

D = density

m = mass

M = Molarity

$1 \text{ cm}^3 = 1 \text{ mL}$

THERMOCHEMISTRY

$$q = mC\Delta T$$

$$q = mH_f \quad q = mH_v$$

$$\Delta H^\circ = \sum \Delta H_f^\circ(\text{products}) - \sum \Delta H_f^\circ(\text{reactants})$$

$$C \text{ of } H_2O(s) = 2.03 \frac{J}{g \cdot ^\circ C}$$

$$C \text{ of } H_2O(l) = 4.184 \frac{J}{g \cdot ^\circ C}$$

$$C \text{ of } H_2O(g) = 2.06 \frac{J}{g \cdot ^\circ C}$$

q = heat (Joules)

m = mass (grams)

C = specific heat capacity ($\frac{J}{g \cdot ^\circ C}$)

T = temperature ($^\circ C$)

H° = standard enthalpy

ΔH_f° = standard enthalpy of formation


Heat of fusion of water, H_f = 334 J/g

Heat of vaporization of water, H_v = 2257 J/g

SIGNIFICANT FIGURE RULES

1. Non-zero digits and zeros between non-zero digits are always significant.
2. Leading zeros are not significant.
3. Trailing zeros are significant if a decimal point is shown.
4. Zeros with a bar over them are significant.
5. In scientific notation, all digits of the coefficient are significant.
6. In a logarithm, there are as many digits after the decimal point as there are significant figures in the original value.

ACTIVITY SERIES FOR METALS



SOLUBILITY RULES FOR SALTS

These rules are written by priority, top to bottom. Rules higher in the list override rules lower in the list. Each statement implies that the listed ion behaves as the title states.

**Examining several sources will yield several slightly different sets of solubility rules. For our work and assessments, these are the only rules to consider.*

Always Soluble

1. alkali metal cations (Li^+ , Na^+ , K^+ , Rb^+ , Cs^+) and NH_4^+
2. NO_3^- , $C_2H_3O_2^-$, NO_2^- , ClO_3^- , ClO_4^-

Generally Soluble

3. Cl^- , Br^- , I^- except when bound to Hg_2^{2+} , Ag^+ , Pb^{2+}
4. SO_4^{2-} except Ca^{2+} , Sr^{2+} , Ba^{2+} , Hg_2^{2+} , Ag^+ , Pb^{2+}

Generally Insoluble

5. F^- , CO_3^{2-} , PO_4^{3-} , S^{2-} , SO_3^{2-} , $C_2O_4^{2-}$, CrO_4^{2-} , $Cr_2O_7^{2-}$, CN^- except Rule 1.
6. O^{2-} , OH^- except Rule 1. Oxides and hydroxides of Ca^{2+} , Sr^{2+} , Ba^{2+} are slightly soluble (this is still considered insoluble)